



WIPES

ONE STEP[†] DISINFECTANT CLEANER & DEODORIZER
For Veterinary[§] Use

Reference Sheet

KEEP OUT OF REACH
OF CHILDREN

Manufactured by:

Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

EPA Reg. No. 74559-10
EPA Est. No. 74559-CAN-1
EPA Est. No. 075272-WI-001



This product is a one-step[†] disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous surfaces.

It cleans quickly by removing common soils found in animal housing facilities, veterinary clinics, animal shelters and kennels, pet boarding facilities, grooming facilities, and pet stores.

This product is designed for use on hard, non-porous surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile and stainless steel associated with high touch surfaces, animal equipment, animal crates, cages, carts, examination tables, grooming equipment, reception areas, veterinary care surfaces or any hard, non-porous washable surfaces where disinfection is required. It also reduces odor-causing bacteria, leaving surfaces smelling clean and fresh.

*VIRUCIDAL: 1 MINUTE, UNLESS OTHERWISE NOTED

Adenovirus Type 8
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Avian Reovirus
Bovine Viral Diarrhea Virus
Canine Distemper Virus
Canine Influenza A Virus (H3N2)
Canine Parvovirus – 5 minutes
Cytomegalovirus
Duck Hepatitis B Virus
Equine Influenza A Virus (H3N8) (EIV)
Feline Calicivirus
Feline Herpes Virus (Feline Rhinotracheitis)
Feline Infectious Peritonitis Virus (FIPV)
Feline Leukemia Virus (FeLV)
Feline Panleukopenia Virus
Hepatitis B Virus (Duck Hepatitis B Virus as the surrogate)
Hepatitis C Virus (Bovine Viral Diarrhea Virus as the surrogate)
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2
HIV Type 1 Strain HTLV III_B
Human Coronavirus
Influenza A Virus
Influenza B Virus
Newcastle Disease Virus
Norovirus (Feline Calicivirus as the surrogate)
Pandemic Influenza A Virus H1N1
Parainfluenza Virus Type 3
Pseudorabies Virus
Respiratory Syncytial Virus (RSV)
Rhinovirus Type-37
Rotavirus (Strain WA)
SARS-Related Coronavirus 2 (Human Coronavirus SARS-CoV-2 (the virus that causes COVID-19[§]))

Vaccinia Virus

Meets surface disinfection recommendations under the OSHA Bloodborne Pathogen Standard for *HBV, *HCV & *HIV-1

BACTERICIDAL: 1 MINUTE

Acinetobacter baumannii (Multi-Drug Resistant)
Bordetella bronchiseptica
Bordetella pertussis
Burkholderia cepacia
Enterobacter aerogenes
Enterobacter cloacae (New Delhi metallo-beta-lactamase 1 (NDM -1))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Escherichia coli
Escherichia coli O157:H7
Escherichia coli (extended spectrum beta-lactamase (ESBL))
Escherichia coli (New Delhi metallo-beta-lactamase 1 (NDM-1))
Klebsiella pneumoniae
Klebsiella pneumoniae (extended spectrum beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-beta-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Legionella pneumoniae
Listeria monocytogenes
Pasteurella multocida
Pseudomonas aeruginosa
Pseudomonas aeruginosa (metallo-beta-lactamase (MBL) positive)
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus
Staphylococcus aureus Community Acquired Methicillin Resistant Genotype USA 400 (CA-MRSA 400)
Staphylococcus aureus (Methicillin Resistant (MRSA))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Staphylococcus pseudintermedius
Stenotrophomonas (Pseudomonas) maltophilia
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Streptococcus pyogenes

FUNGICIDAL: 30 SECONDS, UNLESS OTHERWISE NOTED

Candida albicans – 1 minute
Microsporum canis – 1 minute
Trichophyton interdigitale

APPROVED

By Junette Tabadero at 4:13 pm, May 29, 2025

REF226 (24/355)

MOLD/MILDEW (MILDEWCIDAL ACTIVITY): 4 MINUTES

Aspergillus brasiliensis (formerly known as *Aspergillus niger*)

TUBERCULOCIDAL: 1 MINUTE

Mycobacterium bovis (BCG)

NON-FOOD CONTACT SANITIZER: 10 SECONDS

Acinetobacter baumannii

Campylobacter jejuni

Enterobacter aerogenes

Enterococcus faecalis (Vancomycin Resistant (VRE))

Escherichia coli O157:H7

Listeria monocytogenes

Pseudomonas aeruginosa

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus

Staphylococcus aureus (Methicillin Resistant (MRSA))

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

***KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of (inanimate) hard, non-porous surfaces / objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain visibly wet for 30 seconds to kill HIV-1, HBV and HCV.

Disposal of Infectious Material: Cleaning materials used that may contain blood / body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

For Use as a One-Step Cleaner/Disinfectant

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous surfaces.
3. Allow surface to remain visibly wet for 1 minute.[§]
4. Allow to air dry.

For Use in Animal Housing Facilities such as Veterinary Clinics and Animal Hospitals

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter from floors, walls and surfaces of facilities occupied by animals.

3. Empty all feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Pull towelette from canister and wipe hard, non-porous surfaces.
5. All surfaces must remain visibly wet for 1 minute.[§]
6. Allow to air dry.
7. Thoroughly scrub all treated automatic feeders and waterers with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/or Deodorizer

1. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
2. Allow to air dry.

For Use as a Non-Food Contact Sanitizer

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain visibly wet for 10 seconds.
4. Wipe surface dry or allow to air dry.

To Kill Mold and Mildew

1. Pre-clean surfaces.
2. Pull towelette from canister and wipe product onto hard, non-porous surfaces until completely wet.
3. Allow the surface to remain visibly wet for 4 minutes.
4. Allow to air dry.

Refill Pouch:

1. Remove lid from canister.
2. Place saturated wipes in the empty canister.
3. Use scissors to cut the refill pouch.
4. Pull out corner of first wipe from center of roll and push through "X" opening on lid.
5. Replace lid and pull wipe at an angle. When finished, close lid to retain moisture.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in a cool dry area, away from direct sunlight and heat. Do not contaminate water, food or feed by storage and disposal.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush. Tightly reseal canister, bucket, or soft pack between uses to retain moisture. Do not reuse or refill except as described in the directions for use. Refill only with this product. Offer empty container for recycling. If recycling is not available, discard in trash.

DISCLAIMER: You must read and follow the instructions on the label for our product and use the product for the particular applications specified on the product label. To the full extent permitted under applicable law, Virox Technologies Inc. shall not be liable for any damages whatsoever, arising out of or otherwise in connection with any failure to follow those instructions or any use of its products which is not specified on the product's label. The instructions on the product label are not a warranty, express or implied, and Virox Technologies, Inc. expressly disclaims any implied warranties, including any warranty of fitness for a particular purpose. Under applicable law, some of the above waivers or disclaimers of warranty, or exclusions of liability for certain types of damages may not apply to you. In such cases, our liability will be limited to the fullest extent permitted by applicable law.

[~]Accelerated Hydrogen Peroxide[®] and Design are trademarks of Diversey Inc.
[~]"Accelerated" does not imply increased product efficacy.

[†] When used according to disinfection use directions.

[§] Use a 5 minute contact time against Canine Parvovirus.

[§] Hard, non-porous surfaces